

Contents lists available at [SciVerse ScienceDirect](http://SciVerse.ScienceDirect.com)

Discrete Mathematics

journal homepage: www.elsevier.com/locate/disc

A sufficient condition for Kim's conjecture on the competition numbers of graphs

Akira Kamibepu

Interdisciplinary Faculty of Science and Engineering, Shimane University, Shimane 690-8504, Japan

ARTICLE INFO

Article history:

Received 9 September 2010

Received in revised form 22 August 2011

Accepted 28 November 2011

Available online 23 December 2011

Keywords:

Competition graph

Competition number

Hole

ABSTRACT

A *hole* of a graph G is an induced cycle of length at least 4. Kim (2005) [3] conjectured that the competition number $k(G)$ is bounded by $h(G) + 1$ for any graph G , where $h(G)$ is the number of holes of G . Li and Chang (2009) [5] proved that the conjecture is true for a graph whose holes all satisfy a property called 'independence'. In this paper, by using similar proof techniques in Li and Chang (2009) [5], we prove the conjecture for graphs satisfying two conditions that allow the holes to overlap a lot.

© 2011 Elsevier B.V. All rights reserved.

1. Introduction and preliminaries

Throughout the present paper, all graphs and digraphs (directed graphs) are finite and simple. We use $V(G)$ and $V(D)$ for the vertex set of a graph G or a digraph D . We use $E(G)$ for the edge set of a graph G . An edge with endpoints u and v is denoted by uv . We use $A(D)$ for the set of directed edges of a digraph D . Each element (u, v) of $A(D)$ is called an *arc* from u to v . A digraph is *acyclic* if it contains no directed cycles. The *competition graph* of a digraph D (see [7] for its background) is the graph $C(D)$ on $V(D)$ defined by

$$E(C(D)) = \{uv \mid \text{there is a vertex } x \in V(D) \text{ such that } (u, x), (v, x) \in A(D)\}.$$

For graphs G_1 and G_2 , $G_1 \cup G_2$ is the graph defined by $V(G_1 \cup G_2) = V(G_1) \cup V(G_2)$ and $E(G_1 \cup G_2) = E(G_1) \cup E(G_2)$. Let G be a graph and I_k a set of k isolated vertices, each of which is not a vertex of G , where k is a positive integer. Define I_0 to be the null graph, with no vertices. It is not difficult to see that there is an acyclic digraph D on $V(G) \cup I_{|E(G)|}$ such that $C(D) = G \cup I_{|E(G)|}$. The *competition number* of G , denoted $k(G)$, is defined by

$$k(G) = \min\{k \mid \text{there is an acyclic digraph } D \text{ such that } C(D) = G \cup I_k\}.$$

A *hole* of a graph G is an induced cycle of length at least 4. Researchers have observed a relation between the competition number $k(G)$ of G and the number $h(G)$ of holes of G . Roberts [7] proved that $k(G) \leq 1$ holds for a chordal graph G , that is, for a graph without holes (see Theorem 2.7). Cho and Kim [1] proved that $k(G) \leq 2$ holds for a graph G with exactly one hole. Kim [3] proposed the following conjecture.

Conjecture 1.1 (Kim [3], 2005). *The competition number of a graph G is at most $h(G) + 1$.*

Li and Chang [6] proved that $k(G) \leq 3$ holds for a graph with exactly two holes. In terms of arrangement of holes, they studied a relation between the competition number of a graph and the number of holes of the graph. A hole C of G is *independent* if C satisfies the following conditions with respect to any other hole C' of G :

E-mail address: kamibepu@riko.shimane-u.ac.jp.




Fig. 1. A hole-simple graph that is not pairwise edge-disjoint and has a non-independent hole.




Fig. 2. A graph without Condition (2), whose holes all are independent and pairwise edge-disjoint.

- (i) C and C' have at most two common vertices,
- (ii) if C and C' have two common vertices, then they have one common edge, and
- (iii) if there is a hole C' such that (ii) holds, then C has length at least 5.

Li and Chang [5] showed that $k(G) \leq h(G) + 1$ holds for a graph G whose holes all are independent. For a hole C of G , let X_C be the set of vertices adjacent to all vertices of C . It is not difficult to see that X_C is a clique for an independent hole C . Hence,

Condition (1) X_C is a clique of G for each hole C of G

holds when all holes are independent.

Other conditions on a graph G under which $k(G) \leq h(G) + 1$ holds were also given in [2,4]. The author [2] presented the following condition on graphs:

- for each hole C of a graph, there is an edge which is contained only in C among all induced cycles of the graph.

Kim et al. [4] considered graphs whose holes all are pairwise edge-disjoint. Such graphs are called *hole-edge-disjoint*. The above conditions on graphs imply that any distinct two holes overlap at most one edge.

We present a condition on graphs that allows the holes to overlap a lot. In this paper, a *walk* W from u to v means that u and v are not internal vertices of W . For a hole C of a graph G , a walk W from u to v is said to be *C -avoiding* if internal vertices of W are not in $V(C) \cup X_C$. For a hole C of G and $uv \in E(C)$, let

$$S_{C,uv} = \{x \in V(G) \mid x \text{ is an internal vertex of a } C\text{-avoiding walk from } u \text{ to } v\}, \quad \text{and}$$

$$T_{C,uv} = \{x \in V(G) \mid x \text{ is an internal vertex of a non-}C\text{-avoiding walk from } u \text{ to } v\}.$$

We consider the following condition for a graph G :

Condition (2) $S_{C,uv} \cap T_{C,uv} = \emptyset$ for any hole C of G and any edge $uv \in E(C)$.

A graph is *hole-simple* if it satisfies Conditions (1) and (2). By using similar proof techniques in [5], we prove that $k(G) \leq h(G) + 1$ holds for hole-simple graphs. Hole-simple graphs may have holes that overlap a lot, violating the conditions required in [2,4,5].

The following two examples show that the condition in [4] that all holes are pairwise edge-disjoint, the condition in [5] that all holes are independent, and Conditions (1) and (2) do not imply each other.

Example 1.2. The graph in Fig. 1 is hole-simple, but it is not hole-edge-disjoint. It has a hole that is not independent. We can find two holes of length 6 that overlap at 4 edges.

Example 1.3. The graph in Fig. 2 is not hole-simple. It does not satisfy Condition (2), even though all holes are independent and pairwise edge-disjoint. Actually, for the edge uv of the hole C in the graph, the path (u, y, v) is C -avoiding, so the vertex y is in $S_{C,uv}$. Note that y is also in $T_{C,uv}$, because the walk (u, y, x, y, v) is non- C -avoiding.

In the rest of this section, we recall some basic notions on graphs. Let G be a graph. For a subset S of $V(G)$, the subgraph of G induced by S , denoted $G[S]$, is the subgraph with vertex set S whose edges consist of all edges of G joining two vertices of S . A subset X of $V(G)$ is a *clique* of G if X is a nonempty subset of $V(G)$ such that $G[X]$ is complete. For $u, v \in V(G)$ and $uv \notin E(G)$, let $G + uv$ be the graph with $V(G + uv) = V(G)$ and $E(G + uv) = E(G) \cup \{uv\}$. For $uv \in E(G)$, let $G - uv$ be the graph with $V(G - uv) = V(G)$ and $E(G - uv) = E(G) \setminus \{uv\}$. For $S \subseteq V(G)$, let $G - S = G[V(G) \setminus S]$. A subset $S \subset V(G)$ is called a *vertex cut* of G if $G - S$ has more components than G . Two paths from u to v are *disjoint* if u and v are their only common vertices.

2. The competition numbers of hole-simple graphs

For any clique Q of G , Li and Chang [5] investigated induced subgraphs G_1 and G_2 of G such that

- $G = G_1 \cup G_2$,
- $V(G_1) \cap V(G_2) = \{u, v\} \cup X_C$ is a clique in G , where C is a hole of G_1 and uv is an edge of C ,
- Q is a clique in $G_1 - uv$, and
- both $G_1 - uv$ and G_2 have at most $h(G) - 1$ holes.

They constructed such induced subgraphs for graphs whose holes all are independent. We also construct such induced subgraphs for hole-simple graphs. Unless otherwise specified, in this section we consider a hole-simple graph G . For a hole C of G and $uv \in E(C)$, let $G_1 = G - S_{C,uv}$ and $G_2 = G[\{u, v\} \cup X_C \cup S_{C,uv}]$. Note that $V(G_1) \cap V(G_2) = \{u, v\} \cup X_C$. Under Condition (1), this set is a clique in G .

Lemma 2.1. For a hole C in a graph G , no vertex of $S_{C,uv}$ is adjacent to a vertex of $V(G) \setminus (V(C) \cup X_C \cup S_{C,uv})$.

Proof. If there are vertices $x \in S_{C,uv}$ and $y \in V(G) \setminus (V(C) \cup X_C \cup S_{C,uv})$ such that $xy \in E(G)$, then there is a C -avoiding walk W from u to v including x . Let W_1 from u to x and W_2 from x to v be subwalks of W . The walk $W_1 y W_2$ is also a C -avoiding walk from u to v . Hence, $y \in S_{C,uv}$, a contradiction. \square

Lemma 2.2. For a hole C in a graph G satisfying Condition (2), no vertex of $S_{C,uv}$ is adjacent to a vertex of $V(C) \setminus \{u, v\}$.

Proof. If there are vertices $x \in S_{C,uv}$ and $y \in V(C) \setminus \{u, v\}$ such that $xy \in E(G)$, then there is a C -avoiding walk W from u to v containing x . Let W_1 from u to x and W_2 from x to v be subwalks of W . The walk $W_1 y W_2$ is non- C -avoiding, and hence $x \in T_{C,uv}$. This contradicts Condition (2). \square

Lemmas 2.1 and 2.2 imply that $V(G_1) \cap V(G_2) = \{u, v\} \cup X_C$ is a vertex cut of G if $S_{C,uv} \neq \emptyset$. Hence, $G = G_1 \cup G_2$.

Remark 2.3. By the same argument as in the proof of Lemma 2.2, Condition (2) implies that no vertex of $S_{C,uv}$ is adjacent to a vertex of X_C , so no vertex of $S_{C,uv}$ is adjacent to a vertex of $X_C \cup V(C) \setminus \{u, v\}$. Thus, no vertex of $S_{C,uv}$ are adjacent to a vertex of $V(G) \setminus (\{u, v\} \cup S_{C,uv})$ under Condition (2). Note that $\{u, v\}$ is also a vertex cut of G if $S_{C,uv} \neq \emptyset$.

Lemma 2.4. Let G be a graph with Condition (2) and C a hole of G . Let e and e' be edges of $E(C)$ such that $e \cap e' = \emptyset$. For any $x \in e \cup S_{C,e}$ and any $y \in e' \cup S_{C,e'}$, x is not adjacent to y in G unless $xy \in E(C)$. Also we have $(e \cup S_{C,e}) \cap (e' \cup S_{C,e'}) = \emptyset$.

Proof. Note that $e' \cup S_{C,e'} \subset T_{C,e} \subset V(G) \setminus (e \cup S_{C,e})$ by Condition (2). Hence, $x \in S_{C,e}$ is not adjacent to $y \in e' \cup S_{C,e'}$ by Remark 2.3. We also reach contradictions in other cases by the same argument as above. \square

Lemma 2.5. Let G be a graph satisfying Condition (2). For any hole C and any edge $uv \in E(C)$, no pair (P_1, P_2) of disjoint paths from u to v satisfies the following conditions:

- P_i is a non- C -avoiding path for $i \in \{1, 2\}$ and
- $P_i + uv$ is a hole of G for $i \in \{1, 2\}$.

Proof. Otherwise, for a hole C and $uv \in E(C)$, let P_1 and P_2 be two such paths from u to v . Note that P_i does not have vertices of X_C for $i \in \{1, 2\}$. If P_i has a vertex w_i of X_C , w_i is not in $\{u, v\}$ since $X_C \cap V(C) = \emptyset$. Now, either $w_i u$ or $w_i v$ is a chord of the hole $C_i = P_i + uv$ of G . Hence, P_i must have vertices of $V(C) \setminus \{u, v\}$ for $i \in \{1, 2\}$. Let x_i be a vertex of $V(C) \setminus \{u, v\}$ such that $x_i \in V(P_i)$ for $i \in \{1, 2\}$ (see Fig. 3). Now, P_2 is a C_1 -avoiding path. Otherwise, P_2 must have a vertex z of X_{C_1} since P_1 and P_2 are disjoint. Now, either zu or zv is a chord of C_2 , a contradiction.

Since we can take the subpaths P from x_1 to x_2 and P^{-1} from x_2 to x_1 of C without uv , the walk $P_2^u P^{-1} P_2^v$ is a non- C_1 -avoiding walk having vertices of $S_{C_1,uv}$, where P_2^u from u to x_2 and P_2^v from x_2 to v are the subpaths of P_2 . This is contrary to Condition (2). \square

Lemma 2.6. Let G be a hole-simple graph and C a hole of G and $uv \in E(C)$. If $S_{C,uv} = \emptyset$, the graph $G - uv$ is also a hole-simple graph that satisfies $h(G - uv) < h(G)$.




Fig. 3. Disjoint paths P_1 and P_2 from u to v , each of which is non- C -avoiding.

Proof. By the assumption $S_{C,uv} = \emptyset$, every walk from u to v is non- C -avoiding except for uv . We consider the family $\mathcal{P} = \mathcal{P}_1 \cup \mathcal{P}_2$ of all non- C -avoiding paths from u to v , where

$\mathcal{P}_1 = \{P \mid P \text{ is a non-}C\text{-avoiding path from } u \text{ to } v \text{ such that } P + uv \text{ is a hole of } G\},$ and

$\mathcal{P}_2 = \{P \mid P \text{ is a non-}C\text{-avoiding path from } u \text{ to } v \text{ such that } P + uv \text{ is not a hole of } G\}.$

The path P satisfying $P + uv = C$ is in \mathcal{P}_1 . Note that a path in \mathcal{P}_2 does not form a hole of $G - uv$ with any other paths from u to v . Actually, for a non- C -avoiding path (u, x, v) , the vertex x must be in X_C that is a clique in G by Condition (1). Hence, only pairs of disjoint paths in \mathcal{P}_1 may form holes of $G - uv$ that are not holes of G . By Lemma 2.5, all holes of $G - uv$ are holes of G . Thus, we have

$$h(G - uv) = h(G) - |\mathcal{P}_1| \leq h(G) - 1 < h(G).$$

The graph $G - uv$ satisfies Condition (2) since all holes of $G - uv$ are holes of G . We prove that $G - uv$ also satisfies Condition (1). Suppose to the contrary that there is a hole C' of $G - uv$ such that $\{u, v\} \subseteq X_{C'}$ holds. Note that $V(C) \cap X_C = \emptyset$ for each hole C .

If $V(C') \cap (V(C) \cup X_C) = \emptyset$, the path (u, x, v) is C -avoiding in G for any $x \in V(C')$, and hence $x \in S_{C,uv}$. This contradicts the assumption $S_{C,uv} = \emptyset$.

If $V(C') \cap (V(C) \cup X_C) \neq \emptyset$, let x be a vertex of $V(C') \cap (V(C) \cup X_C)$. Note that $V(C) \cap V(C') = \emptyset$. If not, we have a vertex $y \in V(C') \cap (V(C) \setminus \{u, v\})$. Then, either uy or vy is a chord of C in G , a contradiction. Thus, the vertex x is in $V(C') \cap X_C$. Note that the number of vertices of $V(C') \cap X_C$ is at most 2. Since C' is a cycle of length at least 4, there is a vertex $z \in V(C') \setminus X_C$. Hence, $z \in V(C')$ and $z \notin V(C) \cup X_C$ since $V(C) \cap V(C') = \emptyset$. Now, the path (u, z, v) is a C -avoiding in G , and hence $z \in S_{C,uv}$. This also contradicts the assumption $S_{C,uv} = \emptyset$.

Thus, $X_{C'}$ does not include $\{u, v\}$ for each hole C' of $G - uv$. This implies that $G - uv$ satisfies Condition (1). \square

We need the following useful result to show our main theorem.

Theorem 2.7 (Roberts [7]). Let G be a chordal graph and Q a clique of G . There is an acyclic digraph D such that $C(D) = G \cup I_1$ and all vertices of Q have only outgoing arcs of D .

Theorem 2.8. Let G be a hole-simple graph and Q a clique of G . There is an acyclic digraph D such that $C(D) = G \cup I_{h(G)+1}$ and all vertices of Q have only outgoing arcs in D . Consequently, $k(G) \leq h(G) + 1$.

Proof. We prove this theorem by induction on the number of holes of a graph. If G is a graph without holes, that is, G is a chordal graph, the theorem is true for G by Theorem 2.7.

Let G be a hole-simple graph with at least one hole. Assume that the theorem is true for a hole-simple graph with at most $h(G) - 1$ holes.

Let C be a hole of G and e an edge of C . If $(e \cup S_{C,e}) \cap Q \neq \emptyset$, we may assume that $(e' \cup S_{C,e'}) \cap Q = \emptyset$ for $e' \in E(C)$ such that $e \cap e' = \emptyset$ by Lemma 2.4. Since C is a cycle of length at least 4, there is an edge uv of C such that $\{u, v\} \cup S_{C,uv} \cap Q = \emptyset$. Let $G_1 = G - S_{C,uv}$ and $G_2 = G[\{u, v\} \cup X_C \cup S_{C,uv}]$. By Lemmas 2.1 and 2.2 and Condition (1), the set $V(G_1) \cap V(G_2) = \{u, v\} \cup X_C$ is a clique vertex cut of G if $S_{C,uv} \neq \emptyset$. Hence, $G = G_1 \cup G_2$ and $h(G_2) = h(G) - h(G_1)$ since any hole of G belongs to either G_1 or G_2 . Also note that both G_1 and G_2 are hole-simple.

Note that $h(G_2) < h(G)$ since G_2 does not include the hole C . Then, by the induction hypothesis, there is an acyclic digraph D_2 such that $C(D_2) = G_2 \cup I_{h(G_2)+1}$ and all vertices of the clique $\{u, v\} \cup X_C$ have only outgoing arcs in D_2 . Note that C is a hole of G_1 , Q is a clique of G_1 , and $S_{C,uv} = \emptyset$ in G_1 . By Lemma 2.6, the graph $G_1 - uv$ is hole-simple, and $h(G_1 - uv) < h(G_1) \leq h(G)$. By the induction hypothesis, there is an acyclic digraph D_1 such that $C(D_1) = (G_1 - uv) \cup I_{h(G_1-uv)+1}$ and all vertices of the clique Q have only outgoing arcs in D_1 . Now, consider the digraph D' with $V(D') = V(D_1) \cup V(D_2)$ and $A(D') = A(D_1) \cup A(D_2)$. Note that D' is an acyclic digraph such that $C(D') = G \cup I_{h(G_1-uv)+h(G_2)+2}$ and all vertices of Q have only outgoing arcs in D' , where $h(G_1 - uv) + h(G_2) \leq h(G_1) - 1 + h(G_2) = h(G) - 1$. Thus, $D = D' \cup I_{h(G_1-1)-h(G_1-uv)}$ is a desired acyclic digraph. \square

Remark 2.9. In the proof of Theorem 2.8, we may let $G_2 = G[\{u, v\} \cup S_{C,uv}]$ instead of letting $G_2 = G[\{u, v\} \cup X_C \cup S_{C,uv}]$ since $\{u, v\}$ is also a vertex cut of G under Condition (2) if $S_{C,uv} \neq \emptyset$ (see Remark 2.3).

Acknowledgments

The author would like to thank anonymous referees and an editor for their careful reading of the manuscript and many constructive suggestions and comments on this paper.

References

- [1] H.H. Cho, S.-R. Kim, The competition number of a graph having exactly one hole, *Discrete Math.* 303 (2005) 32–41.
- [2] A. Kamibeppu, An upper bound for the competition numbers of graphs, *Discrete Appl. Math.* 158 (2010) 154–157.
- [3] S.-R. Kim, Graphs with one hole and competition number one, *J. Korean Math. Soc.* 42 (6) (2005) 1251–1264.
- [4] S.-R. Kim, J.Y. Lee, Y. Sano, The competition number of a graph whose holes do not overlap much, *Discrete Appl. Math.* 158 (2010) 1456–1460.
- [5] B.-J. Li, G.J. Chang, The competition number of a graph with exactly h holes, all of which are independent, *Discrete Appl. Math.* 157 (2009) 1337–1341.
- [6] B.-J. Li, G.J. Chang, The competition number of a graph with exactly two holes, *J. Comb. Optim.* (2010), doi:10.1007/s10878-010-9331-9 (Available online first from May 7).
- [7] F.S. Roberts, Food webs, competition graphs, and the boxicity of ecological phase space, in: Y. Alavi, D. Lick (Eds.), *Theory and Applications of Graphs*, in: *Lecture Notes in Mathematics*, vol. 642, Springer-Verlag, 1978, pp. 447–490.